


INFLAMMATORY BOWEL DISEASES

Introduction

#2

- Inflammatory bowel disease (IBD) represents a group of intestinal disorders that cause prolonged inflammation of the digestive tract.
- It is a spectrum of chronic idiopathic inflammatory condition.

Colon



Cecum
portion
of large
intestine



Inflammatory
bowel
disease (IBD)

Ileum
portion
of small
intestine


Classification

Ulcerative colitis: Ulcerative colitis is a disease that causes mucosal inflammation and sores (ulcers) in the lining of the large intestine (colon).

Chron's disease: Crohn's disease is a chronic, relapsing and remitting inflammatory disease of the gastrointestinal tract, affecting any site from mouth to anus.

Crohn's Disease

Distribution



Ulcerative Colitis

Colonic Complications




Inflammatory Bowel Disease

Healthy

Crohn's disease

Ulcerative colitis



Healthy Colon

Ulcerative Colitis

Crohn's Disease

Etiology

Infectious agents

Viruses (Measles)

Bacteria (Mycobacteria)

Genetics

Environmental factors

Diet

Smoking

Psychological factors


Stress

Emotional or physical trauma

Pathophysiology

ALTERED MUCOSAL IMMUNE RESPONSE


- Dietary and bacterial antigens penetrate into the intestinal wall and activates the immune system.
- This causes increased production of pro-inflammatory mediators which will lead to inflammation of the mucosal layer.



Pathophysiology


8

EPITHELIAL DEFECTS



Cont.

- Variety of epithelial defects have been described in Crohn's disease and Ulcerative Colitis.
- Defects in epithelial cells will lead to influx of bacterial components such as dendritic cells and macrophages which activates CD+4 cells.
- Activated CD+4 cells activate other inflammatory cells like B-cells and variable T-cells or recruit more inflammatory cells by stimulation of homing receptor on leucocytes and vascular epithelium.
- Inflammation in IBD is maintained by an influx of leukocytes from the vascular system into sites of active disease. This influx is promoted by expression of adhesion molecules (such as α 4-integrins) on the surface of endothelial cells in the microvasculature in the area of inflammation.



Clinical Manifestations

Clinical symptoms are same in both case.

- Diarrhoea
- Abdominal pain, cramping & bloating due to bowel obstruction
- Hematochezia : Blood in stool
- Low fever
- Decreased appetite
- Weight loss and anorexia
- Fatigue
- Arthritis

Diagnosis

- Physical Examination
- Endoscopy
- Biopsy
- Radiology
- Blood Test

Physical Examination

#13


- The main features to look for are: oral aphthosis, abdominal tenderness and masses, anal tags, fissure and fistulae, nutritional deficiency.
- An important feature in children is growth retardation.



Endoscopy/ Colonoscopy/ Sigmoidoscopy

14

- Colonoscopy helps to determine the pattern and severity of colonic and terminal ileum inflammation and allows biopsies to be obtained.
- Endoscopic features are aphthous ulcers, deeper ulceration, postinflammatory polyps (which indicate previous severe inflammation), but always accompanied by intervening normal mucosa, which is an important differential feature between CD and UC .



Biopsy

15

- Rectal and colonic biopsies should be examined to find the nature of the inflammation (ulcerative colitis versus CD), collagenous colitis or microscopic inflammation if macroscopic appearance is normal, and infection.



Radiology

Barium enema

- Barium inserted into rectum
- Fluoroscopy used to image bowel
- Rarely used due to colonoscopy
- Useful for identifying colonic strictures or colonic fistulae

Barium Small bowel follow-through X-ray

- Barium sulfate suspension drink
- Fluoroscopic images of bowel taken over time
- Useful for looking for inflammation and narrowing of small bowel

Blood Test

- Anemia may be present due to blood loss (iron deficiency), chronic inflammation or B12 malabsorption (macrocytic).
- Hypoalbuminemia suggests severe disease with denutrition. The best markers of inflammation severity are elevation of the C-reactive protein and platelet count.
- Anti-saccharomyces cerevisiae antibodies (ASCA) are positive in 50-60% of CD patients while anti-neutrophil polynuclear antibodies (ANCA) are positive in 50-60% of UC patients.

Inflammatory Bowel Disease: Treatment

