



19 MAT 201

TRANSFORMS AND PARTIAL DIFFERENTIAL EQUATIONS

UNIT - I - FOURIER SERIES

BASIC FORMULAS:

- ① $\frac{d}{dx} (x^n) = nx^{n-1}$
- ② $\frac{d}{dx} (e^{ax}) = ae^{ax}$
- ③ $\frac{d}{dx} (\sin ax) = a \cos ax$
- ④ $\frac{d}{dx} (\cos ax) = -a \sin ax$
- ⑤ $\int \sin ax \, dx = -\frac{\cos ax}{a} + c$
- ⑥ $\int \cos ax \, dx = \frac{\sin ax}{a} + c$
- ⑦ $\int e^{ax} \, dx = \frac{e^{ax}}{a} + c$
- ⑧ $\int x^n \, dx = \frac{x^{n+1}}{n+1} + c$
- ⑨ $\int (ax+b)^n \, dx = \frac{(ax+b)^{n+1}}{a(n+1)} + c$
- ⑩ $\int \frac{dx}{x} = \log x + c$



Bernoulli's formula:

$$* \int uv \, dx = uv_1 - u'v_2 + u''v_3 - \dots$$

$$* \int e^{ax} \sin bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx)$$

$$* \int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx)$$

$$* \sin(A+B) + \sin(A-B) = 2 \sin A \cos B$$

$$* \sin(A+B) - \sin(A-B) = 2 \cos A \sin B$$

$$* \cos(A+B) + \cos(A-B) = 2 \cos A \cos B$$

$$* \cos(A+B) - \cos(A-B) = -2 \sin A \sin B$$

$$* \sin 2x = 2 \sin x \cos x$$

$$* \sin^2 x = \frac{1 - \cos 2x}{2}$$

$$* \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$* \sin n\pi = 0 \text{ for all values of } n, \text{ if } n \text{ is an integer}$$

$$\sin \pi = \sin 2\pi = \sin 3\pi = \dots = \sin n\pi = 0$$

$$* \cos n\pi = (-1)^n, n \text{ is an integer}$$

$$\cos \pi = \cos 3\pi = \cos 5\pi = \cos 7\pi = \dots = (-1)$$

$$\cos 2\pi = \cos 4\pi = \cos 6\pi = \dots = 1$$

$$* \sin(-\theta) = -\sin \theta$$

$$\cos(-\theta) = \cos \theta$$



Problems based on Bernoulli's formula:

① Evaluate $\int x^2 \sin x \, dx$

Soln: $I = \int uv \, dx = u v_1 - u' v_2 + u'' v_3 - \dots$

$u = x^2 \quad v = \sin x$

$u' = 2x \quad v_1 = -\cos x$

$u'' = 2 \quad v_2 = -\sin x$

$u''' = 0 \quad v_3 = \cos x$

$\therefore I = [-x^2 \cos x + 2x \sin x + 2 \cos x] + C$

② Evaluate $\int_0^{2\pi} x^2 \cos nx \, dx$

Soln:

$u = x^2 \quad , \quad v = \cos nx$

$u' = 2x \quad ; \quad v_1 = \frac{\sin nx}{n}$

$u'' = 2 \quad ; \quad v_2 = \frac{-\cos nx}{n^2}$

$u''' = 0 \quad ; \quad v_3 = \frac{-\sin nx}{n^3}$

$\therefore I = \left[x^2 \frac{\sin nx}{n} + 2x \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_0^{2\pi}$

$= 0 + 4\pi \left(\frac{1}{n^2} \right) - 0 - 0$

$I = \frac{4\pi}{n^2}$



3) Evaluate $\int_0^{2\pi} x \sin nx \, dx$

Soln:

$$\begin{array}{l} u = x \\ u' = 1 \\ u'' = 0 \end{array} \quad \begin{array}{l} V = \sin nx \\ V_1 = \frac{-\cos nx}{n} \\ V_2 = \frac{-\sin nx}{n^2} \end{array}$$

$$I = \left[-x \frac{\cos nx}{n} + \frac{\sin nx}{n^2} \right]_0^{2\pi}$$

$$\boxed{I = \frac{-2\pi}{n}}$$

4) Evaluate $\int_{-\pi}^{\pi} x^2 \sin nx \, dx$

Soln:

$$\begin{array}{l} u = x^2 \\ u' = 2x \\ u'' = 2 \\ u''' = 0 \end{array} \quad \begin{array}{l} V = \sin nx \\ V_1 = \frac{-\cos nx}{n} \\ V_2 = \frac{-\sin nx}{n^2} \\ V_3 = \frac{\cos nx}{n^3} \end{array}$$

$$I = \left[-x^2 \frac{\cos nx}{n} + 2x \frac{\sin nx}{n^2} + 2 \frac{\cos nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= -\pi^2 \frac{(-1)^n}{n} + 0 + 2 \frac{\cos n\pi}{n^3} + \pi^2 \frac{(-1)^n}{n} - 0 -$$

$$\boxed{I = 0}$$

$$\frac{2 \cos n\pi}{n^3}$$

$$(\cos(-\theta) = \cos \theta)$$



⑤ Evaluate $\int_0^{2\pi} (x+x^2) \cos nx \, dx$

Soln:

$$u = x + x^2$$

$$v = \cos nx$$

$$u' = 1 + 2x$$

$$v_1 = \frac{\sin nx}{n}$$

$$u'' = 2$$

$$v_2 = -\frac{\cos nx}{n^2}$$

$$u''' = 0$$

$$v_3 = \frac{\sin nx}{n^3}$$

$$I = \left[(x+x^2) \frac{\sin nx}{n} + (1+2x) \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_0^{2\pi}$$

$$= 0 + (1+2\pi) \frac{1}{n^2} + \frac{1}{n^2} - 0 - \frac{1}{n^2}$$

$$I = \frac{2\pi}{n^2}$$

⑥ Evaluate $\int_{-\pi}^{\pi} x \cos nx \, dx$

Soln:

$$u = x$$

$$v = \cos nx$$

$$u' = 1$$

$$v_1 = \frac{\sin nx}{n}$$

$$u'' = 0$$

$$v_2 = -\frac{\cos nx}{n^2}$$

$$I = \left[x \frac{\sin nx}{n} + \frac{\cos nx}{n^2} \right]_{-\pi}^{\pi}$$

$$= 0 + \frac{(-1)^n}{n^2} - 0 - \frac{(-1)^n}{n^2}$$

$$I = 0$$



UNIT - I FOURIER SERIES

Periodic function:

A function $f(x)$ which satisfies the relation $f(x+T) = f(x)$ for all x and some T is called a periodic function. The smallest positive number T for which the relation holds is called the period of $f(x)$.

Example:

- * $\sin x, \cos x$ are periodic function with period 2π .
- * $\sin nx$ and $\cos nx$ are periodic function with period $\frac{2\pi}{n}$.
- * $\tan x$ is a periodic function with period π .

Dirichlet's Conditions:

- * $f(x)$ is periodic, single valued and finite.
- * $f(x)$ has finite number of finite discontinuities in any one period.
- * $f(x)$ has a finite number of maxima or minima.