



## DEPARTMENT OF MATHEMATICS

### UNIT-III PARTIAL DIFFERENTIAL EQUATIONS

#### PARTIAL DIFFERENTIAL EQUATIONS

Defn:

A partial differential equation is an equation involving a function of two or more variables and some of its partial derivatives. Therefore, a partial differential equation contains one dependent and more than one independent variable.

Let  $z = f(x, y)$  is a function of  $x$  &  $y$  then  $z$  is the dependent variable and  $x, y$  are independent variable. The partial derivatives of  $z$  with respect to  $x$  and  $y$  are  $\frac{\partial z}{\partial x}$ ,  $\frac{\partial z}{\partial y}$ ,  $\frac{\partial^2 z}{\partial x^2}$ ,  $\frac{\partial^2 z}{\partial y^2}$ ,  $\frac{\partial^2 z}{\partial x \partial y}$  and we shall use the following notations:

$$\frac{\partial z}{\partial x} = p, \quad \frac{\partial z}{\partial y} = q, \quad \frac{\partial^2 z}{\partial x^2} = r, \quad \frac{\partial^2 z}{\partial y^2} = t, \quad \frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x} = s //$$

#### ORDER OF PDE :

The order of a PDE is the order of a highest partial derivative occurring in the equation.

Eg:  $\frac{\partial u}{\partial x} + \left(\frac{\partial u}{\partial y}\right)^2$ , order is 1, Degree is 2.

Eg:  $\left(\frac{\partial u}{\partial x}\right)^3 + \left(\frac{\partial^2 u}{\partial x^2}\right) + 2\left(\frac{\partial u}{\partial t}\right) = p$ , order is 2, Degree is 1.



## DEPARTMENT OF MATHEMATICS

### UNIT-III PARTIAL DIFFERENTIAL EQUATIONS

1) Form the partial diff. Eqn. by eliminating the arbitrary constants in  $z = ax + by + a^2 + b^2$

$$z = ax + by + a^2 + b^2 \quad \text{--- (1)}$$

p.d.w.r. to  $x$ .

$$\frac{\partial z}{\partial x} = a$$

$$p = a$$

p.d.w.r. to  $y$ .

$$\frac{\partial z}{\partial y} = b$$

$$q = b$$

Sub (a) & (b) in (1)

$$z = px + qy + p^2 + q^2$$

2)  $z = (x^2 + a)(y^2 + b)$

p.d.w.r. to  $x$ .

$$\frac{\partial z}{\partial x} = 2x(y^2 + b)$$

$$p = 2x(y^2 + b)$$

$$y^2 + b = \frac{p}{2x}$$

Arbitrary constant  $\rightarrow$  A.C.

Indep. variable  $\rightarrow$  I.V

A.C  $\leq$  I.V then use p & q

A.C  $>$  I.V then use

P, q, r, s, t



## DEPARTMENT OF MATHEMATICS

### UNIT-III PARTIAL DIFFERENTIAL EQUATIONS

p.d.w.r. to  $y$ .

$$\frac{\partial z}{\partial y} = 2y(x^2 + a)$$

$$q = 2y(x^2 + a)$$

$$\Rightarrow x^2 + a = \frac{q}{2y}$$

$$\therefore \text{Eqn. becomes } z = \frac{p}{2x} \cdot \frac{q}{2y}$$

$$\Rightarrow 4xyz = pq$$

$$) z = (x-a)^2 (y-b)^2$$

p.d.w.r. to  $x$ .

$$\frac{\partial z}{\partial x} = 2(x-a)(y-b)^2$$

$$p = 2(x-a)(y-b)^2$$

$$\frac{\partial^2 z}{\partial x^2} = 2(y-b)^2$$

$$r = 2(y-b)^2 \Rightarrow (y-b)^2 = \frac{r}{2}$$

p.d.w.r. to  $y$ .

$$\frac{\partial z}{\partial y} = 2(y-b)(x-a)^2$$

$$\frac{\partial^2 z}{\partial y^2} = 2(x-a)^2$$

$$t = 2(x-a)^2 \Rightarrow (x-a)^2 = \frac{t}{2}$$

$$\therefore \text{Eqn. becomes } 4z = rt$$



## DEPARTMENT OF MATHEMATICS

### UNIT-III PARTIAL DIFFERENTIAL EQUATIONS

$$1) z = f(x+at) + g(x-at)$$

p.d.w.r. to  $x$

$$\frac{\partial z}{\partial x} = f'(x+at) + g'(x-at)$$

$$p = f'(x+at) + g'(x-at)$$

p.d.w.r. to  $y$

$$q = 0$$

p.d.w.r. to  $x$

$$\frac{\partial^2 z}{\partial x^2} = f''(x+at) + g''(x-at)$$

$$r = f''(x+at) + g''(x-at)$$

p.d.w.r. to  $t$

$$\frac{\partial z}{\partial t} = f'(x+at) \cdot a + g'(x-at) \cdot (-a)$$

$$q_t = f'(x+at) \cdot a - g'(x-at) \cdot a$$

$$\frac{\partial^2 z}{\partial t^2} = f''(x+at) \cdot a^2 + g''(x-at) \cdot a^2$$

$$t = f''(x+at) \cdot a^2 + g''(x-at) \cdot a^2 \therefore$$

$$t = a^2 [f''(x+at) + g''(x-at)]$$

$$\boxed{t = a^2 r}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore – 35



## DEPARTMENT OF MATHEMATICS

### UNIT-III PARTIAL DIFFERENTIAL EQUATIONS