



SNS COLLEGE OF TECHNOLOGY



•

R .SRILEKHA
AP/ CHEMISTRY/ SNSCT

•



ELECTROCHEMICAL SERIES OR EMF SERIES

- ❖ The increasing values of standard reduction potential is called as electrochemical series.
- ❖ The standard electrode potential of a number of electrodes are given in table. The values are determined potentiometrically by combining the electrode with the another standard electrodes, whose electrode potential is zero.

ELECTRODE	ELECTRODE REACTION	E° VOLTS	NATURE
Li+/Li	$\text{Li}^+ + e \rightleftharpoons \text{Li}$	- 3.01	<p style="text-align: center;">Anodic</p> <p style="text-align: center;">↑</p> <p style="text-align: center;">Pt. reference</p> <p style="text-align: center;">↓</p> <p style="text-align: center;">Cathodic</p>
Mg ²⁺ /Mg	$\text{Mg}^{2+} + 2e \rightleftharpoons \text{Mg}$	- 2.37	
Pb ²⁺ / Pb	$\text{Pb}^{2+} + 2e \rightleftharpoons \text{Pb}$	- 1.12	
Zn ²⁺ /Zn	$\text{Zn}^{2+} + 2e \rightleftharpoons \text{Zn}$	- 0.76	
Fe ²⁺ /Fe	$\text{Fe}^{2+} + 2e \rightleftharpoons \text{Fe}$	- 0.44	
Sn ²⁺ /Sn	$\text{Sn}^{2+} + 2e \rightleftharpoons \text{Sn}$	- 0.136	
H ⁺ /H ₂	$2\text{H}^+ + 2e \rightleftharpoons \text{H}_2$	- 0.00	
Cu ²⁺ / Cu	$\text{Cu}^{2+} + 2e \rightleftharpoons \text{Cu}$	+0.34	
Ag ⁺ /Ag	$\text{Ag}^+ + e \rightleftharpoons \text{Ag}$	+0.80	
Au ⁺ / Au	$\text{Au}^+ + e \rightleftharpoons \text{Au}$	+1.50	
Y ₂ F ₂ / F ⁻	$\text{Y}_2\text{F}_2 + e \rightleftharpoons \text{F}^-$	+2.87	



APPLICATION OF ELECTROCHEMICAL SERIES

❖ CALCULATION OF STANDARD EMF OF THE CELL:

The standard emf of the cell (E°) can be calculated if the standard electrode potential values are known using the following reaction.

$$E^\circ_{\text{cell}} = E^\circ_{\text{RHE}} - E^\circ_{\text{LHE}}$$



RELATIVE CASE OF OXIDATION OR REDUCTION



Higher the value of standard reduction potential (+ve value) greater is the tendency to get reduced. (i.e metals on the top (- ve value) are more easily ionized).

- a) The fluorine has higher positive value of standard reduction potential (+ 2.87 V) and shows higher tendency towards reduction.
- b) The lithium has higher negative value (- 3.01 V) and shows higher tendency towards oxidation.



DISPLACEMENT OF ONE ELEMENT BY THE OTHER

Metals which lie higher in the emf series can displace those elements which lie below them in the series.

For example : We may know whether Cu will displace Zn from the solution or vice versa. We know that standard reduction potential of Cu and Zn i.e

$$E^{\circ} \text{Cu}^{2+} / \text{Cu} = + 0.34 \text{ V}$$

$$E^{\circ} \text{Zn}^{2+} / \text{Zn} = - 0.76 \text{ V}$$

So, Cu^{2+} has a great tendency to acquire Cu form than Zn^{2+} has acquiring Zn form



❖ DETERMINATION OF EQUILIBRIUM CONSTANT FOR THE REACTION.

Standard electrode potential can also be used to determine the standard free energy change (ΔG°) and equilibrium constant (K) for the reaction. We know that

$$- \Delta G^\circ = RT \ln k = 2.303 RT \log K$$

$$\log K = - \Delta G^\circ / 2.303 RT$$

$$\log K = nF E^\circ / 2.303 RT \text{ i.e. } \Delta G^\circ = nF E^\circ$$


From the value of E° , the equilibrium constant for the cell reaction can be calculated.




DISPLACEMENT BEHAVIOR OF HYDROGEN



Metals with negative reduction potential will displace the hydrogen from an acid solution.



From the value of E° , the equilibrium constant for the cell reaction can be calculated. The metal with positive reduction potential will not displace the hydrogen from an acid





THANK YOU