

SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution) Coimbatore-641035.



UNIT-III COMPLEX DIFFERENTIATION

Analytic function

Unl17 - 3

Introduction

If x & y are roal numbers then z=x+iy & called a complex number where & is called real part of x y is called imaginary part of z and the value of is JT. The complex number a-in is called the complex conjugate of x and it &s denoted by Z. is) Z = x-14

Noto :

- 1. IZ1 = V22+42
- 2. | Z2 | = ZZ
- 5. ZZ = 22+42= v2
- 4. 121 = 121
- 5. Real past of $z = \frac{z+\overline{z}}{2}$ 6. Imaginary past of $z = \frac{z-\overline{z}}{2}$
- z=rela le alla polar form of z

Function of complex variable

W= f(x)= u(x,y) + PV (x,y) whome u(x,y) & von, y) are real var Pables.

Analytic function a function is said to be analytic at a point if its derivative oxists not only that point but also some neighbourhood of that point



SNS COLLEGE OF TECHNOLOGY



(An Autonomous Institution)
Coimbatore-641035.

UNIT-III COMPLEX DIFFERENTIATION

Analytic function

1. Show that the function $f(x) = \overline{x}$ is nowhow different table.

Garden
$$f(x) = \overline{x} = x - iy$$
 $u + iy = x - iy$
 $\Rightarrow u = x$ and $v = -y$
 $u_x = 1$ $v_x = 0$

$$uy=0$$
 $v_y=-1$

HOME Use # Vy and Uy = - Vze Hence C-R eans we not satisfical.

 \Rightarrow f(x)= \overline{x} is not differentiable anywhere (on) nowhere afferentiable.

2. Determine whether the function axy + i (x2 y2) is analytic or not.

Let
$$f(z) = 2xy + i(x^2 - y^2)$$

 $u + iv = 2xy + i(x^2 - y^2)$

$$\Rightarrow$$
 $u = 2xy$ and $v = x^2 - y^2$
 $u_x = 2x$

$$uy = 2x$$
 $vy = -2y$

$$\Rightarrow$$
 ux \neq Vy and uy \neq -Vx

C.R eans are not satisfied. Hence f(x) is not an analytic function.

3. Let $f(z) = z^3$ be analytic. Justify. Soin.

Let
$$f(z) = z^3$$

 $u + iv = (z + iy)^3$
 $= z^3 + 3z^2(iy) + 3z(iy)^2 + (iy)^3$
 $= z^3 + i3z^2y - 3zy^2 - iy^3$

E



SNS COLLEGE OF TECHNOLOGY



(An Autonomous Institution) Coimbatore-641035.

UNIT-III COMPLEX DIFFERENTIATION

Analytic function

$$u+iv = \begin{bmatrix} x^3 - 3xy^2 \end{bmatrix} + i \begin{bmatrix} 3x^3y - y^3 \end{bmatrix}$$

$$\Rightarrow u = x^3 - 3xy^2 \quad \text{and} \quad v = 3x^3y - y^3$$

$$u_x = 3x^3 - 3y^2 \quad v_x = 6xy$$

$$u_y = -6xy \quad v_y = -3y^3 + 3x^3$$

$$\Rightarrow u_x = v_y \quad \text{and} \quad u_y = -v_x$$

$$CR \quad \text{cqns} \quad \text{and} \quad u_y = -v_x$$

$$CR \quad \text{cqns} \quad \text{and} \quad \text{gatisfied.}$$
Hence, $f(x)$ is analytic.

A. FROD the constants a, b, c 9, f(x) = setay+i(bx+ le analytec. Soln.

Let
$$f(x) = x + ay + i(bx + cy)$$

 $u + iv = x + ay + i(bx + cy)$
Here $u = x + ay$ and $v = bx + cy$
 $u = 1$ $v = b$
 $u = a$ $v = c$

Since
$$f(x)$$
 is analytic.
 $\Rightarrow ux = Vy \text{ and } uy = -Vx$
 $1 = C$
 $\alpha = -b$
 $\alpha = -b$