



Fourier Series

Some Basic Formulas :

1.  $\int \sin x dx = -\cos x$
2.  $\int \cos x dx = \sin x$
3.  $\sin 0 = 0$
4.  $\sin \frac{\pi}{2} = 1$
5.  $\sin n\pi = 0 ; \sin (n+1)2\pi = 0 ; \sin (n+1)\pi = 0$
6.  $\cos 0 = 1$
7.  $\cos \frac{\pi}{2} = 0$
8.  $\cos n\pi = (-1)^n$
9.  $\cos (n+1)2\pi = 1$
10.  $\sin A \cos B = \frac{1}{2} [\sin(A+B) + \sin(A-B)]$
11.  $\cos A \sin B = \frac{1}{2} [\sin(A+B) - \sin(A-B)]$
12.  $\cos A \cos B = \frac{1}{2} [\cos(A+B) + \cos(A-B)]$
13.  $\sin A \sin B = \frac{1}{2} [\cos(A-B) - \cos(A+B)]$
14. Bernoulli's formula :

$$\int u v dx = u v_1 - u' v_2 + u'' v_3 - u''' v_4 + \dots$$

15.  $\int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2+b^2} [a \cos bx + b \sin bx]$
16.  $\int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2+b^2} [a \sin bx - b \cos bx]$



Periodic function :

A function  $f(x)$  is said to be periodic with period ' $p$ ', if for all  $x$ ,  $f(x+p) = f(x)$

where  $p$  is a positive constant, the least value of  $p$  which is called the period of  $f(x)$ .

Eg:  $f(x) = \sin x = \sin(x+2\pi) = \sin(x+4\pi) = \dots$

So,  $\sin x$  is a period of  $2\pi$

Darboux's condition :

Any function  $f(x)$  can be developed as Fourier Series  $\frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$

where  $a_0, a_n$  and  $b_n$  are constants, provided

- i).  $f(x)$  is periodic, single valued and finite.
- ii).  $f(x)$  has a finite no. of finite discontinuities and no infinite discontinuity.
- iii).  $f(x)$  has atmost a finite number of maxima and minima.

Fourier series :

A function  $f(x)$  is periodic and satisfies Darboux's conditions, then it can be represented by an infinite series is called the Fourier series as

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

where  $a_0, a_n$  and  $b_n$  are Fourier coefficients.



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

## FOURIER SERIES- PROBLEMS ON (0,2π)

Euler's Formula:

If a function  $f(x)$  defined in  $c < x < c+2\pi$  can be expanded as the infinite trigonometric series,

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

where  $a_0 = \frac{1}{\pi} \int_c^{c+2\pi} f(x) dx$

$$a_n = \frac{1}{\pi} \int_c^{c+2\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_c^{c+2\pi} f(x) \sin nx dx$$

Problems based on Bernoulli's formula:

i). Find  $\int_0^{2\pi} x \sin x dx$ .

Soln.:

$$\int u v dx = u v - u' v_1 + u'' v_2 - u''' v_3 + \dots$$

Now,  $\int_0^{2\pi} x \sin x dx$

$$= \left[ x(-\cos x) - 1(-\sin x) + 0 \right]_0^{2\pi}$$

$$= [-x \cos x + \sin x]_0^{2\pi}$$

$$= [(0 \cos 2\pi + \sin 2\pi) - 0]$$

$$\int_0^{2\pi} x \sin x dx = -2\pi$$

$$\begin{cases} u = x & v = \sin x \\ u' = 1 & v_1 = -\cos x \\ u'' = 0 & v_2 = -\sin x \end{cases}$$

$$\therefore \cos 2\pi = 1 \\ \sin 2\pi = 0$$

ii). Evaluate  $\int_{-\pi}^{\pi} (x+x^2) \cos nx dx$

Soln.:

$$\begin{cases} u = x + x^2 & v = \cos nx \\ u' = 1+2x & v_1 = \frac{\sin nx}{n} \\ u'' = 2 & v_2 = -\frac{\cos nx}{n^2} \\ u''' = 0 & v_3 = -\frac{\sin nx}{n^3} \end{cases}$$



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON (0,2π)

$$\begin{aligned}
 \int uv dx &= u_1 - u' v_2 + u'' v_3 - u''' v_4 + \dots \\
 \int_{-\pi}^{\pi} (x+x^2) \cos nx dx &= \left[ (x+x^2) \frac{\sin nx}{n} - (1+2x) \left( -\frac{\cos nx}{n^2} \right) + 2 \left( -\frac{\sin nx}{n^3} \right) - 0 \right]_{-\pi}^{\pi} \\
 &= \left[ (x+x^2) \frac{\sin nx}{n} + (1+2x) \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_{-\pi}^{\pi} \\
 &= \left[ (0+(1+2\pi)) \frac{\cos n\pi}{n^2} - 0 \right] - \left[ (0+(1-2\pi)) \frac{\cos(-n\pi)}{n^2} - 0 \right] \\
 &= (1+2\pi) \frac{(-1)^n}{n^2} - (1-2\pi) \frac{(-1)^n}{n^2} \quad \cos n\pi = (-1)^n \\
 &= (1+2\pi - 1+2\pi) \frac{(-1)^n}{n^2} \quad \cos(-n\pi) = \cos n\pi \\
 &= 4\pi \frac{(-1)^n}{n^2} \quad = (-1)^n
 \end{aligned}$$

Q.  $\int_0^{2\pi} x^2 \cos nx dx$



Problems on (0, 2π)

Formula :

$$\begin{aligned}
 f(x) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \\
 a_0 &= \frac{1}{\pi} \int_0^{2\pi} f(x) dx \quad a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx \\
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx
 \end{aligned}$$

Q. Determine the Fourier series for  $f(x) = x^2$ ,  $(0, 2\pi)$

Soln. :

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \quad \hookrightarrow (1)$$



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON (0,2π)

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x^2 dx$$

$$= \frac{1}{\pi} \left[ \frac{x^3}{3} \right]_0^{2\pi} = \frac{1}{3\pi} [8\pi^3 - 0]$$

$$a_0 = \frac{8}{3} \pi^2$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x^2 \cos nx dx$$

$$\int uv dx = uv_1 - u'v_2 + u''v_3 - \dots$$

$$\begin{aligned} u &= x^2 & v &= \cos nx \\ u' &= 2x & v_1 &= \sin nx/n \\ u'' &= 2 & v_2 &= -\cos nx/n^2 \\ u''' &= 0 & v_3 &= -\sin nx/n^3 \end{aligned}$$

$$= \frac{1}{\pi} \left[ x^2 \frac{\sin nx}{n} - 2x \left( -\frac{\cos nx}{n^2} \right) + 2 \left( -\frac{\sin nx}{n^3} \right) - 0 \right]_0^{2\pi}$$

$$= \frac{1}{\pi} \left[ \frac{x^2 \sin nx}{n} + 2x \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_0^{2\pi}$$

$$= \frac{1}{\pi} \left[ \left( 0 + \frac{4\pi \cdot 0}{n^2} \right) - (0) \right] \quad \because \sin 2n\pi = 0 \\ \cos 2n\pi = 1 \\ \sin 0 = 0$$

$$a_n = \frac{4}{n^2}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x^2 \sin nx dx$$

$$\int uv dx = uv_1 - u'v_2 + u''v_3 - \dots$$

$$\begin{aligned} u &= x^2 & v &= \sin nx \\ u' &= 2x & v_1 &= -\cos nx/n \\ u'' &= 2 & v_2 &= -\sin nx/n^2 \\ u''' &= 0 & v_3 &= \cos nx/n^3 \end{aligned}$$



$$\begin{aligned}
 &= \frac{1}{\pi} \left[ x^2 \left( -\frac{\cos nx}{n} \right) - 2x \left( -\frac{\sin nx}{n^2} \right) + 2 \left( \frac{\cos nx}{n^3} \right) - 0 \right]_0^{2\pi} \\
 &= \frac{1}{\pi} \left[ -x^2 \frac{\cos nx}{n} + 2x \frac{\sin nx}{n^2} + 2 \frac{\cos nx}{n^3} \right]_0^{2\pi} \\
 &= \frac{1}{\pi} \left[ \left( -\frac{4\pi^2}{n} + 0 + \frac{2}{n^3} \right) - \left( 0 + 0 + \frac{2}{n^3} \right) \right] \\
 &= \frac{1}{\pi} \left[ -\frac{4\pi^2}{n} + \frac{2}{n^3} - \frac{2}{n^3} \right]
 \end{aligned}$$

$\therefore \cos 2n\pi = 1$   
 $\sin 2n\pi = 0$   
 $\cos 0 = 1$

$$b_n = -\frac{4\pi}{n}$$

$$\begin{aligned}
 \therefore (1) \Rightarrow f(x) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[ \frac{4\pi}{n^2} \cos nx - \frac{4\pi}{n} \sin nx \right] \\
 &= \frac{4}{3}\pi^2 + 4 \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx - 4\pi \sum_{n=1}^{\infty} \frac{1}{n} \sin nx
 \end{aligned}$$

Q.If  $f(x) = \left(\frac{\pi-x}{2}\right)^2$ ,  $(0, 2\pi)$ , determine fourier series for the function  $f(x)$ .

Sol:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \rightarrow (1)$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \left(\frac{\pi-x}{2}\right)^2 dx$$

$$= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 dx$$

$$= \frac{1}{4\pi} \left[ \frac{(\pi-x)^3}{-3} \right]_0^{2\pi} = -\frac{1}{12\pi} [-\pi^3 - \pi^3]$$

$$= -\frac{1}{12\pi} (-2\pi^3)$$

$$a_0 = \frac{\pi^3}{6}$$



$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} \left(\frac{\pi-x}{2}\right)^2 \cos nx dx \\
 &= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 \cos nx dx \quad \int uv dx = uv_1 - u'v_2 + u''v_3 - \dots \\
 &\qquad\qquad\qquad u = (\pi-x)^2 \quad v = \cos nx \\
 &\qquad\qquad\qquad u' = -2(\pi-x) \quad v_1 = \sin nx/n \\
 &\qquad\qquad\qquad u'' = -2(-1)=2 \quad v_2 = -\cos nx/n^2 \\
 &\qquad\qquad\qquad u''' = 0 \quad v_3 = -\sin nx/n^3 \\
 &= \frac{1}{4\pi} \left[ (\pi-x)^2 \left( \frac{\sin nx}{n} \right) - (-2(\pi-x)) \left( -\frac{\cos nx}{n^2} \right) \right. \\
 &\qquad\qquad\qquad \left. + 2 \left( -\frac{\sin nx}{n^3} \right) - 0 \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[ (\pi-2\pi)^2 \frac{\sin n\pi}{n} - 2(\pi-2\pi) \left( \frac{\cos n\pi}{n^2} \right) - 2 \frac{\sin n\pi}{n^3} \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[ (0 - \frac{2(-\pi)}{n^2} - 0) - (0 - \frac{2\pi}{n^2} - 0) \right] \quad \because \sin 2n\pi = 0 \\
 &\qquad\qquad\qquad \cos 2n\pi = 1 \\
 &\qquad\qquad\qquad \cos 0 = 1 \\
 &= \frac{1}{4\pi} \left[ \frac{2\pi}{n^2} + \frac{2\pi}{n^2} \right] \\
 &= \frac{1}{4\pi} \left[ \frac{4\pi}{n^2} \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{n^2} \\
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} \left(\frac{\pi-x}{2}\right)^2 \sin nx dx \quad \int uv dx = uv_1 - u'v_2 + u''v_3 - \dots \\
 &= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 \sin nx dx \quad u = \left(\frac{\pi-x}{2}\right)^2 \quad v = \sin nx \\
 &\qquad\qquad\qquad u' = -2(\pi-x) \quad v_1 = -\cos nx/n \\
 &\qquad\qquad\qquad u'' = -2(-1)=2 \quad v_2 = -\sin nx/n^2 \\
 &\qquad\qquad\qquad u''' = 0 \quad v_3 = \cos nx/n^3
 \end{aligned}$$



$$\begin{aligned}
 &= \frac{1}{4\pi} \left[ (\pi - x)^2 \left( \frac{\cos nx}{n} \right) - (-2(\pi - x)) \left( -\frac{\sin nx}{n^2} \right) \right. \\
 &\quad \left. + 2 \left( \frac{\cos nx}{n^3} \right) - 0 \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[ -(\pi - x)^2 \frac{\cos nx}{n} - 2(\pi - x) \frac{\sin nx}{n^2} + 2 \frac{\cos nx}{n^3} \right]_0^{2\pi} \\
 &= \frac{1}{4\pi} \left[ \left( -\frac{\pi^2}{n} + 2\pi(0) + \frac{2}{n^3} \right) - \left( -\frac{\pi^2}{n} - 0 + \frac{2}{n^3} \right) \right] \\
 &= \frac{1}{4\pi} \left[ -\frac{\pi^2}{n} + \frac{2}{n^3} + \frac{\pi^2}{n} - \frac{2}{n^3} \right]
 \end{aligned}$$

$$b_n = 0$$

$$\begin{aligned}
 \therefore (i) \Rightarrow f(x) &= \frac{\pi^2/6}{2} + \sum_{n=1}^{\infty} \left[ \frac{1}{n^2} \cos nx + 0 \right] \\
 &= \frac{\pi^2}{12} + \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx
 \end{aligned}$$

3]. Find the fourier series for  $f(x) = \begin{cases} x, & 0 < x < \pi \\ 2\pi - x, & \pi < x < 2\pi \end{cases}$

soln.:

$$\begin{aligned}
 f(x) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \rightarrow (i) \\
 a_0 &= \frac{1}{\pi} \int_0^{2\pi} f(x) dx \\
 &= \frac{1}{\pi} \left[ \int_0^{\pi} x dx + \int_{\pi}^{2\pi} (2\pi - x) dx \right] \\
 &= \frac{1}{\pi} \left[ \left( \frac{x^2}{2} \right)_0^{\pi} + \left( 2\pi x - \frac{x^2}{2} \right)_{\pi}^{2\pi} \right] \\
 &= \frac{1}{\pi} \left[ \left( \frac{\pi^2}{2} - 0 \right) + \left( 4\pi^2 - \frac{4\pi^2}{2} \right) - \left( 2\pi^2 - \frac{\pi^2}{2} \right) \right] \\
 &= \frac{1}{\pi} \left[ \frac{\pi^2}{2} + (2\pi^2 - 2\pi^2 + \frac{\pi^2}{2}) \right] = \frac{1}{\pi} \left[ \frac{2\pi^2}{2} \right] \\
 a_0 &= \pi
 \end{aligned}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

## FOURIER SERIES- PROBLEMS ON (0,2π)

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx \\
 &= \frac{1}{\pi} \left[ \int_0^{\pi} x \cos nx dx + \int_{\pi}^{2\pi} (2\pi - x) \cos nx dx \right] \\
 &\quad \int u v dx = uv_1 - u' v_0 + u'' v_2 - u''' v_4 + \dots \\
 u &= x \quad \left| \begin{array}{l} v = \frac{\cos nx}{2\pi - x} \\ v_1 = -\frac{8\pi n \cos nx}{n^2} \\ v_2 = -\frac{\cos nx}{n^2} \end{array} \right| \quad \begin{array}{l} u = 2\pi - x \\ u' = -1 \\ u'' = 0 \end{array} \\
 u' &= 1 \\
 u'' &= 0 \\
 &= \frac{1}{\pi} \left[ \left( x \frac{\sin nx}{n} - 1 \left( -\frac{\cos nx}{n^2} \right) + 0 \right)_0^\pi + \left( (2\pi - x) \frac{\sin nx}{n} + 1 \left( \frac{\cos nx}{n^2} \right) \right)_0^{2\pi} \right] \\
 &= \frac{1}{\pi} \left[ \left( x \frac{\sin nx}{n} + \frac{\cos nx}{n^2} \right)_0^\pi + \left( (2\pi - x) \frac{\sin nx}{n} - \frac{\cos nx}{n^2} \right)_0^{2\pi} \right] \\
 &= \frac{1}{\pi} \left[ \left( \frac{(-1)^n}{n^2} - \frac{1}{n^2} \right) + \left( \frac{-1}{n^2} + \frac{(-1)^n}{n^2} \right) \right] \\
 &= \frac{1}{\pi} \left[ \frac{2(-1)^n}{n^2} - \frac{2}{n^2} \right] = \frac{2}{\pi n^2} [(-1)^n - 1]
 \end{aligned}$$

$$a_n = \frac{2}{\pi n^2} [(-1)^n - 1]$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx \\
 &= \frac{1}{\pi} \left[ \int_0^{\pi} x \sin nx dx + \int_{\pi}^{2\pi} (2\pi - x) \sin nx dx \right] \\
 &= \frac{1}{\pi} \left[ \left\{ x \left( -\frac{\cos nx}{n} \right) - 1 \left( -\frac{\sin nx}{n^2} \right) + 0 \right\}_0^\pi + \left. \begin{array}{l} v = \sin nx \\ v_1 = -\cos nx/n \\ v_2 = -\sin nx/n^2 \end{array} \right\} \right. \\
 &\quad \left. \left\{ (2\pi - x) \left( -\frac{\cos nx}{n} \right) - (-1) \left( -\frac{\sin nx}{n^2} \right) + 0 \right\}_{\pi}^{2\pi} \right] \\
 &= \frac{1}{\pi} \left[ \left( -x \frac{\cos nx}{n} + \frac{\sin nx}{n^2} \right)_0^\pi + \left( -(2\pi - x) \frac{\cos nx}{n} - \frac{\sin nx}{n^2} \right)_\pi^{2\pi} \right] \\
 &= \frac{1}{\pi} \left[ -\pi \frac{(-1)^n}{n} + \frac{\pi (-1)^n}{n} \right] \\
 b_n &= 0
 \end{aligned}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

## FOURIER SERIES- PROBLEMS ON (0, 2π)

$$f(x) = \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{2}{\pi n^2} [(-1)^{n-1}] \cos nx + 0 \\ = \frac{\pi}{2} + \frac{2}{\pi} \sum_{n=1}^{\infty} [(-1)^{n-1}] \cos nx$$

Q. find the Fourier Series for  $f(x) = x \sin nx, (0, 2\pi)$

Soln. :

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

b

$$= \frac{1}{\pi} \int_0^{2\pi} x \sin nx dx$$

$$= \frac{1}{\pi} \left[ x(-\cos nx) - 1(-\sin nx) \right]_0^{2\pi}$$

$$= \frac{1}{\pi} \left[ -x \cos nx + \sin nx \right]_0^{2\pi}$$

$$= \frac{1}{\pi} [-2\pi - 0] = -\frac{2}{\pi}$$

$$a_0 = -\frac{2}{\pi}$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x \sin nx \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} x \cos nx \sin nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \frac{x}{2} [ \sin(n\pi + nx) - \sin(n\pi - nx) ] dx = \frac{1}{2\pi} \int_0^{2\pi} x \sin(n+1)x dx \\ - \int_0^{2\pi} x \sin(n-1)x dx \}$$

$\neq \frac{1}{2\pi}$

$$u = x \quad v = \sin(n+1)x \\ u' = 1 \quad v_1 = -\frac{\cos(n+1)x}{(n+1)} \\ u'' = 0 \quad v_2 = -\frac{\sin(n+1)x}{(n+1)^2}$$

$$v = \sin(n-1)x \\ v_1 = -\frac{\cos(n-1)x}{(n-1)} \\ v_2 = -\frac{\sin(n-1)x}{(n-1)^2}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



## UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

## FOURIER SERIES- PROBLEMS ON (0,2π)

$$\begin{aligned}
 &= \frac{1}{2\pi} \left[ \left( \int_0^{2\pi} \left( \frac{\cos(n+1)x)}{n+1} - \frac{-\sin(n+1)x)}{(n+1)^2} \right) dx \right] \\
 &\quad + \left[ \int_0^{2\pi} \left( \frac{\cos(n-1)x)}{n-1} - \frac{-\sin(n-1)x)}{(n-1)^2} \right) dx \right] \\
 &= \frac{1}{2\pi} \left[ \left( \int_0^{2\pi} \frac{x \cos(n+1)x}{n+1} dx + \frac{\sin(n+1)x}{(n+1)^2} \right) \Big|_0^{2\pi} + \left( \int_0^{2\pi} \frac{-x \cos(n-1)x}{n-1} dx + \frac{\sin(n-1)x}{(n-1)^2} \right) \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \frac{-2\pi}{n+1} + \frac{2\pi}{n-1} \right] \\
 &= \frac{+2\pi}{2\pi} \left[ \frac{n+1 + n-1}{(n+1)(n-1)} \right]
 \end{aligned}$$

$$a_n = \frac{+2\pi}{2\pi}$$

$$a_1 = \frac{1}{\pi} \int_0^{\pi} x \cos x dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x \sin x dx \cos x dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x dx$$

$$= \frac{1}{2\pi} \left[ -\frac{\cos x}{2} \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} -1$$

$$= \frac{1}{2\pi} \int_0^{2\pi} x \sin x dx$$

$$= \frac{1}{2\pi} \left[ x \left( -\frac{\cos 2x}{2} \right) - \left( \frac{\sin 2x}{4} \right) \right]_0^{2\pi}$$

$$= \frac{1}{2\pi} \left[ -\frac{2\pi}{2} \right]$$

$$a_2 = -1$$

$$\cos(n+1)2\pi = 1$$

$$\sin(n+1)2\pi = 0$$

$$\cos(n-1)2\pi = \cos(2\pi) = 1$$

$$\cos 2\pi \cos 2\pi + \sin 2\pi \sin 2\pi = 1$$

$$\sin(n-1)2\pi = \sin(2\pi - 2\pi) = 0$$

$$\sin 2\pi \cos 2\pi - \cos 2\pi \sin 2\pi = 0$$

$$\int u v dx = u v_i - u' v_2 + u'' v_3 - \dots$$

$$\begin{cases} u = x \\ u' = 1 \\ u'' = 0 \end{cases} \quad 
 \begin{cases} v = \sin 2x \\ v_1 = -\cos 2x/2 \\ v_2 = -\sin 2x/4 \end{cases}$$



$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx dx = \frac{1}{\pi} \int_0^{2\pi} x \sin nx \frac{\sin x}{x} dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} \frac{x}{2} [ \cos(n+1)x - \cos(n-1)x ] dx \\
 &= \frac{1}{2\pi} \left[ \int_0^{2\pi} x \cos(n+1)x dx - \int_0^{2\pi} x \cos(n-1)x dx \right] \\
 &= \frac{1}{2\pi} \left[ \left\{ x \frac{\sin(n+1)x}{n+1} - \left( -\frac{\cos(n+1)x}{(n+1)^2} \right) \right\} \Big|_0^{2\pi} \right. \\
 &\quad \left. - \left\{ x \frac{\sin(n-1)x}{n-1} - \left( -\frac{\cos(n-1)x}{(n-1)^2} \right) \right\} \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{\cos(n+1)2\pi}{n+1} + \frac{\cos(n-1)2\pi}{n-1} \right) - \left( \frac{\cos(n+1)0}{n+1} + \frac{\cos(n-1)0}{n-1} \right) \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right) - \left( \frac{1}{(n+1)^2} + \frac{1}{(n-1)^2} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 b_0 &= 0 \\
 b_1 &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin x dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x \left( \frac{1-\cos 2x}{2} \right) dx \\
 &= \frac{1}{2\pi} \left[ \int_0^{2\pi} x dx - \int_0^{2\pi} \frac{x \cos 2x}{2} dx \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{x^2}{2} \right)_0^{2\pi} - \frac{1}{2} \left\{ x \frac{\sin 2x}{2} - \left( -\frac{\cos 2x}{4} \right) \right\} \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \frac{4\pi^2}{2} - \frac{1}{2} \left\{ \frac{1}{4} - \frac{1}{4} \right\} \right]
 \end{aligned}$$

$$b_1 = \pi$$



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON (0,2π)

$$f(x) = \frac{-\alpha}{2} + a_1 \cos x + \sum_{n=2}^{\infty} a_n \cos nx + b_1 \sin x + \sum_{n=2}^{\infty} b_n \sin nx$$
$$= -1 - \frac{1}{2} \cos x + \sum_{n=2}^{\infty} \frac{1}{n^2-1} \cos nx + \pi \sin x + 0$$
$$= -1 - \frac{\cos x}{2} + 2 \sum_{n=2}^{\infty} \frac{1}{n^2-1} \cos nx + \pi \sin x$$

3).  $f(x) = \begin{cases} 1, & 0 < x < \pi \\ 2, & \pi < x < 2\pi \end{cases}$

4).  $f(x) = e^{-x}, 0 < x < 2\pi$

H.W. D.  $f(x) = \frac{x}{\pi-x}$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON  $(0, 2\pi)$

Method of Variation of Parameters

The second order linear differential eqn is

$$\frac{d^2y}{dx^2} + P \frac{dy}{dx} + Q = X \text{ where } X \text{ is a fn. of } x.$$

CF =  $c_1 f_1 + c_2 f_2$ , where  $c_1, c_2$  are constants  
 $f_1, f_2$  are functions of  $x$ .

$$PI = Pf_1 + Qf_2$$

$$\text{where } P = - \int \frac{f_2 X}{f_1 f_2' - f_1' f_2} dx$$

$$Q = \int \frac{f_2 X}{f_1 f_2' - f_1' f_2} dx$$

$$\int \tan x dx \\ = +\log (\sec x)$$

$$\int \cot x dx \\ = \log (\sin x)$$

$$\int \csc x dx \\ = -\log [\csc x + \cot x]$$

$$\int \sec x dx \\ = \log (\sec x + \tan x)$$

Ex. Solve  $\frac{d^2y}{dx^2} + 4y = 4 \tan 2x$  using method of variation of parameters.

Soln.

$$\text{Given } (D^2 + 4)y = 4 \tan 2x \quad \text{where } x = 4 \tan 2x$$

AE

$$D^2 + 4 = 0$$

$$D^2 = -4$$

$$D = \pm 2i$$

$$CF = C_1 \cos 2x + C_2 \sin 2x$$

$$PI = Pf_1 + Qf_2$$

Here  $f_1 = \cos 2x$

$$f_1' = -2 \sin 2x \quad \left| \begin{array}{l} f_2 = \sin 2x \\ f_2' = 2 \cos 2x \end{array} \right.$$

$$\text{Now } \omega = f_1 f_2' - f_2 f_1'$$

$$= \cos 2x [2 \cos 2x] - \sin 2x (-2 \sin 2x)$$

$$= 2 \cos^2 2x + 2 \sin^2 2x$$

$$= 2 [\cos^2 2x + \sin^2 2x]$$

$$= 2(1) = 2$$



Scanned with  
CamScanner



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON (0,2π)

$$\begin{aligned}
 P &= - \int \frac{f_2 x}{\omega} dx \\
 &= - \int \frac{\sin 2x + \tan 2x}{2} dx \\
 &= -2 \int \sin 2x \frac{\sin 2x}{\cos 2x} dx \\
 &= -2 \int \frac{\sin^2 2x}{\cos 2x} dx \\
 &= -2 \int \frac{1 - \cos^2 2x}{\cos 2x} dx = -2 \left[ \int \frac{1}{\cos 2x} dx - \int \cos 2x dx \right] \\
 &= -2 \left[ \int \sec 2x dx - \int \cos 2x dx \right] \\
 &= -2 \left[ \frac{\log (\sec 2x + \tan 2x)}{2} + \frac{\sin 2x}{2} \right] \\
 P &= -\log (\sec 2x + \tan 2x) + \sin 2x \\
 Q &= \int \frac{f_1 x}{\omega} dx \\
 &= \int \frac{\cos 2x + \tan 2x}{2} dx \\
 &= 2 \int \cos 2x \frac{\sin 2x}{\cos 2x} dx \\
 &= 2 \int \sin 2x dx = 2 \left[ -\frac{\cos 2x}{2} \right] \\
 Q &= -\cos 2x \\
 PI &= Pf_1 + Q f_2 \\
 &= \left[ \log (\sec 2x + \tan 2x) + \sin 2x \right] \cos 2x \\
 &\quad - \cos 2x \sin 2x \\
 PI &= -\log (\sec 2x + \tan 2x) \cos 2x \\
 \therefore y &= CF + PI = C_1 \cos 2x + C_2 \sin 2x - \frac{\log (\sec 2x + \tan 2x)}{\cos 2x}
 \end{aligned}$$



Scanned with  
CamScanner