



4. Test for goodness of fit :

From the sample data we fit a curve or a distribution (binomial, exponential, poisson or normal). We set the null hypothesis as the given sample is from a specified population.

Let O_i and E_i be the observed and expected frequencies such that $\sum O_i = \sum E_i = N$ and the test-statistic is,

$$\chi^2 = \sum_{i=1}^n \frac{(O_i - E_i)^2}{E_i}$$

with $v = n - 1$ d.o.f.

Note :

1. In the case of fitting a Binomial distribution
d.o.f = $n - 1$, for poisson distribution, $v = n - 2$
and for normal distribution $v = n - 3$.

2. If $\chi^2 = 0$, all observed and expected frequencies coincide.



DEPARTMENT OF MATHEMATICS

Problems :

(165)

- ① Fit a binomial distribution for the following data and also test the goodness of fit.

x	0	1	2	3	4	5	6	Total
f	5	18	28	12	7	6	4	80

Solution :

Null hypothesis : H_0 : Binomial fit is a good fit to the data.

Alternative hypothesis : H_1 : Binomial fit is not a good fit to the data.

Calculation of expected frequencies :

x	0	1	2	3	4	5	6	Total
f	5	18	28	12	7	6	4	80
fx	0	18	56	36	28	30	24	192

$$\bar{x} = \frac{\sum fx}{\sum f}$$

$$\bar{x} = \frac{192}{80} = 2.4$$

$$\boxed{\bar{x} = 2.4}$$

For Binomial distribution, mean = np

$$\text{i.e., } \bar{x} = np \Rightarrow np = 2.4$$



DEPARTMENT OF MATHEMATICS

$$6p = 2.4$$

$$\underline{p = 0.4}$$

$$\underline{q = 0.6}$$

The expected frequencies are calculated using the formula, $N \times n C_r p^r q^{n-r}$

where $N = 80$, $n = 6$, $p = 0.4$, $q = 0.6$ &

$$r = 0, 1, 2, \dots, 6$$

$$E(0) = 80 \times 6 C_0 (0.4)^0 (0.6)^6 = 3.73$$

$$E(1) = 80 \times 6 C_1 (0.4)^1 (0.6)^5 = 14.93$$

$$E(2) = 80 \times 6 C_2 (0.4)^2 (0.6)^4 = 24.88$$

$$E(3) = 80 \times 6 C_3 (0.4)^3 (0.6)^3 = 22.12$$

$$E(4) = 80 \times 6 C_4 (0.4)^4 (0.6)^2 = 11.06$$

$$E(5) = 80 \times 6 C_5 (0.4)^5 (0.6) = 2.95$$

$$E(6) = 80 \times 6 C_6 (0.4)^6 (0.6)^0 = 0.33$$

Hence the observed and expected frequencies are given by,

O_i	5	18	28	12	7	6	4
E_i	4	15	25	22	11	3	0



The first class is combined with the second and the last two classes are combined with the last but second class in order to make the expected frequency in each class greater than or equal to 10. Thus, after regrouping, we have,

O_i	23	28	12	17
E_i	19	25	22	14

O_i	E_i	$O_i - E_i$	$(O_i - E_i)^2$	$\frac{(O_i - E_i)^2}{E_i}$
23	19	4	16	0.842
28	25	3	9	0.36
12	22	-10	100	4.545
17	14	3	9	0.643
	80			6.39

Test-Statistic :

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$

$$= 6.39$$

$$\boxed{\chi^2 = 6.39}$$



DEPARTMENT OF MATHEMATICS

Table Value :

At $\alpha = 5\%$ LOS, $\nu = n - k = 4 - 2 = 2$
d.o.f, the table value of χ^2 is given by,

$$\chi_{\alpha}^2 = 5.99$$

Decision :

Since $\chi^2 > \chi_{\alpha}^2$, H_0 is rejected. \therefore The binomial fit is not a good fit for the data.

- ② Fit a poisson distribution for the following distribution and also test the goodness of fit.

x	0	1	2	3	4	5	Total
f	142	156	69	27	5	1	400

Solution :

Null hypothesis : H_0 : Poisson fit is a good fit to the data.

Alternative hypothesis : H_1 : poisson fit is not a good fit to the data.

Calculation of expected frequencies :



DEPARTMENT OF MATHEMATICS

(167)

x	0	1	2	3	4	5	Total
f	142	156	69	27	5	1	400
fx	0	156	138	81	20	5	400

$$\bar{x} = \frac{\sum fx}{\sum f} = \frac{400}{400} = 1$$

For poisson distribution, mean = λ

$$\text{i.e., } \bar{x} = \lambda \Rightarrow \underline{\lambda = 1}$$

The expected frequencies are given by,

$$\frac{N e^{-\lambda} \lambda^r}{r!} \quad \text{where } N = 400, \lambda = 1, r = 0, 1, \dots, 5$$

$$E(0) = \frac{400 \times e^{-1}}{0!} = 147.2$$

$$E(1) = \frac{400 \times e^{-1}}{1!} = 147.2$$

$$E(2) = \frac{400 \times e^{-1}}{2!} = 73.58$$

$$E(3) = \frac{400 \times e^{-1}}{3!} = 24.53$$

$$E(4) = \frac{400 \times e^{-1}}{4!} = 6.13$$

$$E(5) = \frac{400 \times e^{-1}}{5!} = 1.23$$



DEPARTMENT OF MATHEMATICS

Thus,

O_i	142	156	69	27	5	1
E_i	147	147	74	25	6	1

Let us combine the last 3 groups into one, so that $E_i \geq 10$.

O_i	E_i	$O_i - E_i$	$(O_i - E_i)^2$	$(O_i - E_i)^2 / E_i$
142	147	-5	25	0.17
156	147	9	81	0.55
69	74	-5	25	0.34
33	32	1	1	0.03
	400			1.09

Test-Statistic :

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}$$

$$\boxed{\chi^2 = 1.09}$$

Table value :

At $\alpha = 5\%$ LOS, $v = n - 2 = 4 - 2 = 2$

d.o.f, the value of χ^2 in tables is,

$$\boxed{\chi^2_{\alpha} = 5.99}$$

Decision : Since $\chi^2 < \chi^2_{\alpha}$, H_0 is accepted \therefore

Poisson fit is a good fit to the data.



Problems :

- (i) Find if there is any association between extravagance in fathers and extravagance in sons from the following data.

	Extravagant father	Miserly father
Extravagant son	327 ^a	741 ^b
Miserly Son	545 ^c	234 ^d

Determine the coefficient of association also.

Solution :

Null hypothesis : H_0 : The attributes are independent.

Alternative hypothesis : H_1 : The attributes are dependent.

Test-Statistic :

For a 2×2 Contingency table,

$$\chi^2 = \frac{(ad - bc)^2 (a + b + c + d)}{(a + b)(c + d)(a + c)(b + d)}$$

$$= \frac{[(327)(234) - (545)(741)]^2 \times (327 + 234 + 545 + 741)}{(875)(975)(1068)(779)}$$



DEPARTMENT OF MATHEMATICS

$$\chi^2 = 230.24$$

Table value :

At $\alpha = 5\%$ LOS, $v = (p-1)(r-1)$
 $= (2-1)(2-1) = 1$ d.o.f, the table value of χ^2
is given by,

$$\chi_{\alpha}^2 = 3.841$$

Decision:

Since $\chi^2 > \chi_{\alpha}^2$, H_0 is rejected. \therefore The attributes
are dependent. Hence there is an association between
extravagance in fathers and extravagance in sons.

Coefficient of attributes :

$$\begin{aligned} \text{Coefficient of association} &= \frac{ad - bc}{ad + bc} \\ &= \frac{327 \times 234 - 741 \times 545}{327 \times 234 + 741 \times 545} \\ &= \frac{-327330}{480363} \end{aligned}$$

$$Q = -0.6814$$