



## Fourier series

Some basic formulas:

1.  $\int \sin x \, dx = -\cos x$

2.  $\int \cos x \, dx = \sin x$

3.  $\sin 0 = 0$

4.  $\sin \frac{\pi}{2} = 1$

5.  $\sin n\pi = 0$  ;  $\sin (n+1)2\pi = 0$  ;  $\sin (n+1)\pi = 0$

6.  $\cos 0 = 1$

7.  $\cos \frac{\pi}{2} = 0$

8.  $\cos n\pi = (-1)^n$

9.  $\cos (n+1)2\pi = 1$

10.  $\sin A \cos B = \frac{1}{2} [\sin (A+B) + \sin (A-B)]$

11.  $\cos A \sin B = \frac{1}{2} [\sin (A+B) - \sin (A-B)]$

12.  $\cos A \cos B = \frac{1}{2} [\cos (A+B) + \cos (A-B)]$

13.  $\sin A \sin B = \frac{1}{2} [\cos (A-B) - \cos (A+B)]$

14. Bernoulli's formula:

$$\int u v \, dx = u v_1 - u' v_2 + u'' v_3 - u''' v_4 + \dots$$

15.  $\int e^{ax} \cos bx \, dx = \frac{e^{ax}}{a^2 + b^2} [a \cos bx + b \sin bx]$

16.  $\int e^{ax} \sin bx \, dx = \frac{e^{ax}}{a^2 + b^2} [a \sin bx - b \cos bx]$



Periodic function :

A function  $f(x)$  is said to be periodic with period 'p', if for all  $x$ ,  $f(x+p) = f(x)$

where p is a positive constant, the least value of  $p > 0$  which is called the period of  $f(x)$ .

Eg:  $f(x) = \sin x = \sin(x+2\pi) = \sin(x+4\pi) = \dots$

So,  $2\pi$  is a period of  $\underline{2\pi}$

Dirichlet's condition :

Any function  $f(x)$  can be developed as Fourier series  $\left[ \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \right]$  provided.

where  $a_0, a_n$  and  $b_n$  are constants, provided.

- i).  $f(x)$  is periodic, single valued and finite.
- ii).  $f(x)$  has a finite no. of finite discontinuities and no infinite discontinuity.
- iii).  $f(x)$  has at most a finite number of maxima and minima.

Fourier series :

A function  $f(x)$  is periodic and satisfies Dirichlet's conditions, then it can be represented by an infinite series is called the Fourier series as

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

where  $a_0, a_n$  and  $b_n$  are Fourier coefficients.



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON  $(0, 2\pi)$

Euler's Formula:

If a function  $f(x)$  defined in  $c < x < c + 2\pi$  can be expanded as the infinite trigonometric series,

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

where  $a_0 = \frac{1}{\pi} \int_c^{c+2\pi} f(x) dx$  ;  $a_n = \frac{1}{\pi} \int_c^{c+2\pi} f(x) \cos nx dx$

$$b_n = \frac{1}{\pi} \int_c^{c+2\pi} f(x) \sin nx dx$$

Problems based on Bernoulli's formula:

1]. Find  $\int_0^{2\pi} x \sin x dx$ .

Soln.:

$$\int uv dx = uv_1 - u'v_2 + u''v_3 - u'''v_4 + \dots$$

now,  $\int_0^{2\pi} x \sin x dx$

$$= [x(-\cos x) - 1(-\sin x) + 0]_0^{2\pi}$$

$$= [-x \cos x + \sin x]_0^{2\pi}$$

$$\int_0^{2\pi} x \sin x dx = [-2\pi \cos 2\pi + \sin 2\pi] - [0] = -2\pi$$

$$\begin{array}{l|l} u = x & v = \sin x \\ u' = 1 & v_1 = -\cos x \\ u'' = 0 & v_2 = -\sin x \end{array}$$

$\therefore \cos 2\pi = 1$   
 $\sin 2\pi = 0$

2]. Evaluate  $\int_{-\pi}^{\pi} (x + x^2) \cos nx dx$

Soln.:

$$\begin{array}{l|l} u = x + x^2 & v = \cos nx \\ u' = 1 + 2x & v_1 = \frac{\sin nx}{n} \\ u'' = 2 & v_2 = \frac{-\cos nx}{n^2} \\ u''' = 0 & v_3 = \frac{-\sin nx}{n^3} \end{array}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON  $(0, 2\pi)$

Now

$$\int uv dx = uv - u'v_2 + u''v_3 - u'''v_4 + \dots$$

$$\int_{-\pi}^{\pi} (x+x^2) \cos nx dx$$

$$= \left[ (x+x^2) \frac{\sin nx}{n} - (1+2x) \left( -\frac{\cos nx}{n^2} \right) + 2 \left( -\frac{\sin nx}{n^3} \right) - 0 \right]_{-\pi}^{\pi}$$

$$= \left[ (x+x^2) \frac{\sin nx}{n} + (1+2x) \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_{-\pi}^{\pi}$$


$$= \left[ (0+(1+2\pi)) \frac{\cos n\pi}{n^2} - 0 \right] - \left[ (0+(1-2\pi)) \frac{\cos(-n\pi)}{n^2} - 0 \right]$$

$$= (1+2\pi) \frac{(-1)^n}{n^2} - (1-2\pi) \frac{(-1)^n}{n^2}$$

$$= (1+2\pi-1+2\pi) \frac{(-1)^n}{n^2}$$

$$= 4\pi \frac{(-1)^n}{n^2}$$

$\cos n\pi = (-1)^n$   
 $\cos(-n\pi) = \cos n\pi = (-1)^n$



HW  $\int_0^{2\pi} x^2 \cos nx dx$

Problems on  $(0, 2\pi)$

Formula:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx$$

Q. Determine the Fourier series for  $f(x) = x^2$ ,  $(0, 2\pi)$

Soln.:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \quad \rightarrow (1)$$





# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON (0,2π)

$$\begin{aligned}
 a_0 &= \frac{1}{\pi} \int_0^{2\pi} f(x) dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x^2 dx \\
 &= \frac{1}{\pi} \left[ \frac{x^3}{3} \right]_0^{2\pi} = \frac{1}{3\pi} [8\pi^3 - 0]
 \end{aligned}$$

$$a_0 = \frac{8}{3} \pi^2$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x^2 \cos nx dx
 \end{aligned}$$

$$\int u v dx = u v_1 - u' v_2 + u'' v_3 - \dots$$

$$\begin{array}{l}
 u = x^2 \\
 u' = 2x \\
 u'' = 2 \\
 u''' = 0
 \end{array}
 \left\{
 \begin{array}{l}
 v = \cos nx \\
 v_1 = \sin nx/n \\
 v_2 = -\cos nx/n^2 \\
 v_3 = -\sin nx/n^3
 \end{array}
 \right.$$

$$= \frac{1}{\pi} \left[ x^2 \frac{\sin nx}{n} - 2x \left( -\frac{\cos nx}{n^2} \right) + 2 \left( -\frac{\sin nx}{n^3} \right) - 0 \right]_0^{2\pi}$$

$$= \frac{1}{\pi} \left[ \frac{x^2 \sin nx}{n} + 2x \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_0^{2\pi}$$

$$= \frac{1}{\pi} \left[ \left( 0 + \frac{4\pi(1)}{n^2} - 0 \right) - (0) \right] \quad \begin{array}{l} \because \sin 2n\pi = 0 \\ \cos 2n\pi = 1 \\ \sin 0 = 0 \end{array}$$

$$a_n = \frac{4}{n^2}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x^2 \sin nx dx$$

$$\int u v dx = u v_1 - u' v_2 + u'' v_3 - \dots$$

$$\begin{array}{l}
 u = x^2 \\
 u' = 2x \\
 u'' = 2 \\
 u''' = 0
 \end{array}
 \left\{
 \begin{array}{l}
 v = \sin nx \\
 v_1 = -\cos nx/n \\
 v_2 = -\sin nx/n^2 \\
 v_3 = \cos nx/n^3
 \end{array}
 \right.$$



$$= \frac{1}{\pi} \left[ x^2 \left( -\frac{\cos nx}{n} \right) - 2x \left( -\frac{\sin nx}{n^2} \right) + 2 \left( \frac{\cos nx}{n^3} \right) - 0 \right]_{0}^{2\pi}$$

$$= \frac{1}{\pi} \left[ -x^2 \frac{\cos nx}{n} + 2x \frac{\sin nx}{n^2} + 2 \frac{\cos nx}{n^3} \right]_{0}^{2\pi}$$

$$= \frac{1}{\pi} \left[ \left( -\frac{4\pi^2}{n} + 0 + \frac{2}{n^3} \right) - \left( 0 + 0 + \frac{2}{n^3} \right) \right]$$

$$= \frac{1}{\pi} \left[ -\frac{4\pi^2}{n} + \frac{2}{n^3} - \frac{2}{n^3} \right]$$

$$\begin{aligned} \because \cos 2n\pi &= 1 \\ \sin 2n\pi &= 0 \\ \cos 0 &= 1 \end{aligned}$$

$$b_n = -\frac{4\pi}{n}$$

$$\begin{aligned} \therefore (1) \Rightarrow f(x) &= \frac{8/3 \pi^2}{2} + \sum_{n=1}^{\infty} \left[ \frac{4}{n^2} \cos nx - \frac{4\pi}{n} \sin nx \right] \\ &= \frac{4}{3} \pi^2 + 4 \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx - 4\pi \sum_{n=1}^{\infty} \frac{1}{n} \sin nx \end{aligned}$$

Q. If  $f(x) = \left( \frac{\pi-x}{2} \right)^2$ ,  $(0, 2\pi)$ , determine fourier series for the function  $f(x)$ .

Soln.:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \rightarrow (1)$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \left( \frac{\pi-x}{2} \right)^2 dx$$

$$= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 dx$$

$$= \frac{1}{4\pi} \left[ \frac{(\pi-x)^3}{-3} \right]_0^{2\pi} = -\frac{1}{12\pi} \left[ -\pi^3 - \pi^3 \right]$$

$$= -\frac{1}{12\pi} (-2\pi^3)$$

$$a_0 = \frac{\pi^2}{6}$$



$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \left(\frac{\pi-x}{2}\right)^2 \cos nx \, dx$$

$$= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 \cos nx \, dx \quad \int uv \, dx = u v_1 - u' v_2 + u'' v_3 - \dots$$

$$\begin{array}{l} u = (\pi-x)^2 \\ u' = -2(\pi-x) \\ u'' = -2(-1) = 2 \\ u''' = 0 \end{array} \quad \left. \begin{array}{l} v = \cos nx \\ v_1 = \sin nx/n \\ v_2 = -\cos nx/n^2 \\ v_3 = -\sin nx/n^3 \end{array} \right\} \begin{array}{l} \\ \\ \\ \end{array}$$

$$= \frac{1}{4\pi} \left[ (\pi-x)^2 \left(\frac{\sin nx}{n}\right) - (-2(\pi-x)) \left(-\frac{\cos nx}{n^2}\right) + 2 \left(-\frac{\sin nx}{n^3}\right) - 0 \right]_0^{2\pi}$$

$$= \frac{1}{4\pi} \left[ (\pi-x)^2 \frac{\sin nx}{n} - 2(\pi-x) \left(\frac{\cos nx}{n^2}\right) - 2 \frac{\sin nx}{n^3} \right]_0^{2\pi}$$

$$= \frac{1}{4\pi} \left[ \left(0 - 2 \frac{(-\pi)}{n^2} - 0\right) - \left(0 - \frac{2\pi}{n^2} - 0\right) \right] \quad \begin{array}{l} \because \sin 2n\pi = 0 \\ \cos 2n\pi = 1 \\ \cos 0 = 1 \end{array}$$

$$= \frac{1}{4\pi} \left[ \frac{2\pi}{n^2} + \frac{2\pi}{n^2} \right]$$

$$= \frac{1}{4\pi} \left[ \frac{4\pi}{n^2} \right]$$

$$a_n = \frac{1}{n^2}$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \left(\frac{\pi-x}{2}\right)^2 \sin nx \, dx \quad \int uv \, dx = u v_1 - u' v_2 + u'' v_3 - \dots$$

$$\begin{array}{l} u = \left(\frac{\pi-x}{2}\right)^2 \\ u' = -2(\pi-x) \\ u'' = -2(-1) = 2 \\ u''' = 0 \end{array} \quad \left. \begin{array}{l} v = \sin nx \\ v_1 = -\cos nx/n \\ v_2 = -\sin nx/n^2 \\ v_3 = \cos nx/n^3 \end{array} \right\} \begin{array}{l} \\ \\ \\ \end{array}$$

$$= \frac{1}{4\pi} \int_0^{2\pi} (\pi-x)^2 \sin nx \, dx$$



$$\begin{aligned}
&= \frac{1}{4\pi} \left[ (\pi-x)^2 \left( \frac{-\cos nx}{n} \right) - (-2(\pi-x)) \left( -\frac{\sin nx}{n^2} \right) \right. \\
&\quad \left. + 2 \left( \frac{\cos nx}{n^3} \right) - 0 \right]_{0}^{2\pi} \\
&= \frac{1}{4\pi} \left[ -(\pi-x)^2 \frac{\cos nx}{n} - 2(\pi-x) \frac{\sin nx}{n^2} + 2 \frac{\cos nx}{n^3} \right]_{0}^{2\pi} \\
&= \frac{1}{4\pi} \left[ \left( -\frac{\pi^2}{n} + 2\pi(0) + \frac{2}{n^3} \right) - \left( -\frac{\pi^2}{n} - 0 + \frac{2}{n^3} \right) \right] \\
&= \frac{1}{4\pi} \left[ -\frac{\pi^2}{n} + \frac{2}{n^3} + \frac{\pi^2}{n} - \frac{2}{n^3} \right]
\end{aligned}$$

b

$$b_n = 0$$

$$\begin{aligned}
\therefore (1) \Rightarrow f(x) &= \frac{\pi^2/6}{2} + \sum_{n=1}^{\infty} \left[ \frac{1}{n^2} \cos nx + 0 \right] \\
&= \frac{\pi^2}{12} + \sum_{n=1}^{\infty} \frac{1}{n^2} \cos nx
\end{aligned}$$

3]. Find the Fourier series for  $f(x) = \begin{cases} x, & 0 < x < \pi \\ 2\pi - x, & \pi < x < 2\pi \end{cases}$

Soln.:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx] \rightarrow (1)$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$= \frac{1}{\pi} \left[ \int_0^{\pi} x dx + \int_{\pi}^{2\pi} (2\pi - x) dx \right]$$

$$= \frac{1}{\pi} \left[ \left( \frac{x^2}{2} \right)_0^{\pi} + \left( 2\pi x - \frac{x^2}{2} \right)_{\pi}^{2\pi} \right]$$

$$= \frac{1}{\pi} \left[ \frac{1}{2}(\pi^2 - 0) + \left( 4\pi^2 - \frac{4\pi^2}{2} \right) - \left( 2\pi^2 - \frac{\pi^2}{2} \right) \right]$$

$$= \frac{1}{\pi} \left[ \frac{\pi^2}{2} + \left( 2\pi^2 - 2\pi^2 + \frac{\pi^2}{2} \right) \right] = \frac{1}{\pi} \left[ \frac{2\pi^2}{2} \right]$$

$$a_0 = \pi$$





# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore-641035.



UNIT-IV FOURIER SERIES AND FOURIER TRANSFORM

FOURIER SERIES- PROBLEMS ON  $(0, 2\pi)$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

$$= \frac{1}{\pi} \left[ \int_0^{\pi} x \cos nx \, dx + \int_{\pi}^{2\pi} (2\pi - x) \cos nx \, dx \right]$$

$$\int u v \, dx = u v_1 - u' v_2 + u'' v_3 - u''' v_4 + \dots$$

$$\begin{array}{l} u = x \\ u' = 1 \\ u'' = 0 \end{array} \left| \begin{array}{l} v = \cos nx \\ v_1 = \frac{\sin nx}{n} \\ v_2 = -\frac{\cos nx}{n^2} \end{array} \right. \begin{array}{l} u = 2\pi - x \\ u' = -1 \\ u'' = 0 \end{array}$$

$$= \frac{1}{\pi} \left[ \left( x \frac{\sin nx}{n} - 1 \left( -\frac{\cos nx}{n^2} \right) + 0 \right) \Big|_0^{\pi} + \left( (2\pi - x) \frac{\sin nx}{n} + 1 \left( \frac{\cos nx}{n^2} \right) \right) \Big|_{\pi}^{2\pi} \right]$$

$$= \frac{1}{\pi} \left[ \left( x \frac{\sin nx}{n} + \frac{\cos nx}{n^2} \right) \Big|_0^{\pi} + \left( (2\pi - x) \frac{\sin nx}{n} - \frac{\cos nx}{n^2} \right) \Big|_{\pi}^{2\pi} \right]$$

$$= \frac{1}{\pi} \left[ \left( \frac{(-1)^n}{n^2} - \frac{1}{n^2} \right) + \left( -\frac{1}{n^2} + \frac{(-1)^n}{n^2} \right) \right]$$

$$= \frac{1}{\pi} \left[ \frac{2(-1)^n}{n^2} - \frac{2}{n^2} \right] = \frac{2}{\pi n^2} [(-1)^n - 1]$$

$$a_n = \frac{2}{\pi n^2} [(-1)^n - 1]$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

$$= \frac{1}{\pi} \left[ \int_0^{\pi} x \sin nx \, dx + \int_{\pi}^{2\pi} (2\pi - x) \sin nx \, dx \right]$$

$$= \frac{1}{\pi} \left[ \left\{ x \left( -\frac{\cos nx}{n} \right) - 1 \left( -\frac{\sin nx}{n^2} \right) + 0 \right\} \Big|_0^{\pi} + \left\{ (2\pi - x) \left( -\frac{\cos nx}{n} \right) - (-1) \left( -\frac{\sin nx}{n^2} \right) + 0 \right\} \Big|_{\pi}^{2\pi} \right]$$

$$\begin{array}{l} v = \sin nx \\ v_1 = -\cos nx/n \\ v_2 = -\sin nx/n^2 \end{array}$$

$$= \frac{1}{\pi} \left[ \left( -x \frac{\cos nx}{n} + \frac{\sin nx}{n^2} \right) \Big|_0^{\pi} + \left( -(2\pi - x) \frac{\cos nx}{n} - \frac{\sin nx}{n^2} \right) \Big|_{\pi}^{2\pi} \right]$$

$$= \frac{1}{\pi} \left[ -\frac{\pi(-1)^n}{n} + \frac{\pi(-1)^n}{n} \right]$$

$$b_n = 0$$



$$f(x) = \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{2}{\pi n^2} [(-1)^n - 1] (\cos nx + 0)$$

$$= \frac{\pi}{2} + \frac{2}{\pi} \sum_{n=1}^{\infty} [(-1)^n - 1] \cos nx$$

Q. Find the Fourier Series for  $f(x) = x \sin x, (0, 2\pi)$

Soln.:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos nx + b_n \sin nx]$$

$$a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x \sin x dx$$

$$= \frac{1}{\pi} [x(-\cos x) - 1(-\sin x)]_0^{2\pi}$$

$$= \frac{1}{\pi} [-x \cos x + \sin x]_0^{2\pi}$$

$$= \frac{1}{\pi} [-2\pi - 0] = -2$$

$$\begin{array}{l|l} u = x & v = \sin x \\ u' = 1 & v_1 = -\cos x \\ u'' = 0 & v_2 = -\sin x \end{array}$$

$$a_0 = -2$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos nx dx = \frac{1}{\pi} \int_0^{2\pi} x \cos nx \sin x dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} \frac{x}{2} [\sin(n+1)x - \sin(n-1)x] dx = \frac{1}{2\pi} \int_0^{2\pi} x \sin(n+1)x dx - \int_0^{2\pi} x \sin(n-1)x dx$$

$$\neq \frac{1}{2\pi}$$

$$\begin{array}{l|l|l} u = x & v = \sin(n+1)x & v = \sin(n-1)x \\ u' = 1 & v_1 = \frac{-\cos(n+1)x}{(n+1)} & v_1 = \frac{-\cos(n-1)x}{(n-1)} \\ u'' = 0 & v_2 = \frac{-\sin(n+1)x}{(n+1)^2} & v_2 = \frac{-\sin(n-1)x}{(n-1)^2} \end{array}$$



$$\begin{aligned}
 &= \frac{1}{2\pi} \left[ \left( x \left( \frac{\cos(n+1)x}{n+1} \right) - \left( \frac{-\sin(n+1)x}{(n+1)^2} \right) \right) \Big|_0^{2\pi} - \right. \\
 &\quad \left. \left( x \left( \frac{\cos(n-1)x}{n-1} \right) - \left( \frac{-\sin(n-1)x}{(n-1)^2} \right) \right) \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \left( x \frac{\cos(n+1)x}{n+1} + \frac{\sin(n+1)x}{(n+1)^2} \right) \Big|_0^{2\pi} + \left( -x \frac{\cos(n-1)x}{n-1} + \frac{\sin(n-1)x}{(n-1)^2} \right) \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \frac{-2\pi}{n+1} + \frac{2\pi}{n-1} \right] \\
 &= \frac{+2\pi}{2\pi} \left[ \frac{n+1+n-1}{(n+1)(n-1)} \right]
 \end{aligned}$$

$\cos(n+1)2\pi = 1$   
 $\sin(n+1)2\pi = 0$   
 $\cos(n-1)2\pi = \cos(2n\pi)$   
 $\sin(n-1)2\pi = \sin(2n\pi)$   
 $\cos 2n\pi \cos 2\pi + \sin 2n\pi \sin 2\pi = 1$   
 $\sin(n-1)2\pi = \sin(2n\pi - 2\pi) = \sin 2n\pi \cos 2\pi - \cos 2n\pi \sin 2\pi = 0$

$a_n = \frac{+2}{n^2 - 1}$

$a_1 = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos x \, dx$

$= \frac{1}{\pi} \int_0^{2\pi} x \sin x \cos x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \frac{\sin 2x}{2} \, dx$

$= \frac{1}{2\pi} \int_0^{2\pi} x \sin 2x \, dx$

$= \frac{1}{4\pi} [-1]$

$\int u v \, dx = u v_1 - u' v_2 + u'' v_3 - \dots$   
 $u = x \quad \left| \quad v = \sin 2x \right.$   
 $u' = 1 \quad \left| \quad v_1 = -\cos 2x / 2 \right.$   
 $u'' = 0 \quad \left| \quad v_2 = -\sin 2x / 4 \right.$

$= \frac{1}{2\pi} \left[ x \left( \frac{-\cos 2x}{2} \right) - \left( \frac{\sin 2x}{4} \right) \right]_0^{2\pi}$

$= \frac{1}{2\pi} \left[ -\frac{2\pi}{2} \right]$

$a_1 = -1/2$



$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin nx \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin nx \frac{\sin x \, dx}{\frac{1}{2}} \\
 &= \frac{1}{\pi} \int_0^{2\pi} \frac{x}{2} [\cos(n-1)x - \cos(n+1)x] \, dx \\
 &= \frac{1}{2\pi} \left[ \int_0^{2\pi} x \cos(n-1)x \, dx - \int_0^{2\pi} x \cos(n+1)x \, dx \right] \\
 &= \frac{1}{2\pi} \left[ \left\{ x \frac{\sin(n-1)x}{n-1} - \int_0^{2\pi} \left( -\frac{\cos(n-1)x}{(n-1)^2} \right) \right\} \right. \\
 &\quad \left. - \left\{ x \frac{\sin(n+1)x}{n+1} - \int_0^{2\pi} \left( -\frac{\cos(n+1)x}{(n+1)^2} \right) \right\} \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{x \sin(n-1)x}{n-1} + \frac{\cos(n-1)x}{(n-1)^2} \right) \Big|_0^{2\pi} - \left( \frac{x \sin(n+1)x}{n+1} + \frac{\cos(n+1)x}{(n+1)^2} \right) \Big|_0^{2\pi} \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{1}{(n-1)^2} - \frac{1}{(n-1)^2} \right) - \left( \frac{1}{(n+1)^2} - \frac{1}{(n+1)^2} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 b_n &= 0 \\
 b_1 &= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin x \, dx = \frac{1}{\pi} \int_0^{2\pi} x \sin x \sin x \, dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} x \left( \frac{1 - \cos 2x}{2} \right) \, dx \\
 &= \frac{1}{2\pi} \left[ \int_0^{2\pi} x \, dx - \int_0^{2\pi} \frac{x \cos 2x}{2} \, dx \right] \\
 &= \frac{1}{2\pi} \left[ \left( \frac{x^2}{2} \right) \Big|_0^{2\pi} - \frac{1}{2} \left\{ x \frac{\sin 2x}{2} - \int_0^{2\pi} \left( \frac{\cos 2x}{4} \right) \right\} \right] \\
 &= \frac{1}{2\pi} \left[ \frac{4\pi^2}{2} - \frac{1}{2} \left\{ \frac{1}{4} - \frac{1}{4} \right\} \right]
 \end{aligned}$$

$$b_1 = \pi$$





$x$

$$f(x) = \frac{-2}{2} + a_1 \cos x + \sum_{n=2}^{\infty} a_n \cos nx + b_1 \sin x + \sum_{n=2}^{\infty} b_n \sin nx$$

$$= -1 - \frac{1}{2} \cos x + \sum_{n=2}^{\infty} \frac{2}{n^2-1} \cos nx + \pi \sum_{n=2}^{\infty} \sin nx + 0$$

$$= -1 - \frac{\cos x}{2} + 2 \sum_{n=2}^{\infty} \frac{1}{n^2-1} \cos nx + \pi \sum_{n=2}^{\infty} \sin nx$$

HW. 1)  $f(x) = \frac{x}{\pi-x}$

3)  $f(x) = \begin{cases} 1, & 0 < x < \pi \\ 2, & \pi < x < 2\pi \end{cases}$

4)  $f(x) = e^{-x}, 0 < x < 2\pi$



Method of variation of Parameters

The second order linear differential eqn. is

$$\frac{d^2 y}{dx^2} + P \frac{dy}{dx} + Q = X \text{ where } X \text{ is a fn. of } x.$$

CF =  $C_1 f_1 + C_2 f_2$ , where  $C_1, C_2$  are constants  
 $f_1, f_2$  are functions of  $x$ .

PI =  $P f_1 + Q f_2$

where  $P = - \int \frac{f_2 x}{f_1 f_2' - f_1' f_2} dx$

$Q = \int \frac{f_2 x}{f_1 f_2' - f_1' f_2} dx$

$\int \tan x dx = + \log(\sec x)$

$\int \cot x dx = \log(\sin x)$

$\int \operatorname{cosec} x dx = -\log[\csc x + \cot x]$

$\int \sec x dx = \log(\sec x + \tan x)$

II. Solve  $\frac{d^2 y}{dx^2} + 4y = 4 \tan 2x$  using method of variation of parameters.

Soln.

Given  $(D^2 + 4)y = 4 \tan 2x$  where  $X = 4 \tan 2x$

AE

$$m^2 + 4 = 0$$

$$m^2 = -4$$

$$m = \pm 2i$$

$$CF = C_1 \cos 2x + C_2 \sin 2x$$

$$PI = P f_1 + Q f_2$$

Here  $f_1 = \cos 2x$

$$f_1' = -2 \sin 2x$$

$$f_2 = \sin 2x$$

$$f_2' = 2 \cos 2x$$

Now  $w = f_1 f_2' - f_2 f_1'$

$$= \cos 2x [2 \cos 2x] - \sin 2x (-2 \sin 2x)$$

$$= 2 \cos^2 2x + 2 \sin^2 2x$$

$$= 2 [\cos^2 2x + \sin^2 2x]$$

$$= 2(1) = 2$$





$$\begin{aligned}
 P &= - \int \frac{f_2 x}{\omega} dx \\
 &= - \int \frac{\sin 2x + \tan 2x}{2} dx \\
 &= - \frac{1}{2} \int \sin 2x + \frac{\sin 2x}{\cos 2x} dx \\
 &= - \frac{1}{2} \int \frac{\sin^2 2x}{\cos 2x} dx \\
 &= - \frac{1}{2} \int \frac{1 - \cos^2 2x}{\cos 2x} dx = - \frac{1}{2} \left[ \int \frac{1}{\cos 2x} dx - \int \cos 2x dx \right] \\
 &= - \frac{1}{2} \left[ \int \sec 2x dx - \int \cos 2x dx \right] \\
 &= - \frac{1}{2} \left[ \frac{\log |\sec 2x + \tan 2x|}{2} + \frac{\sin 2x}{2} \right]
 \end{aligned}$$

$$P = - \log |\sec 2x + \tan 2x| + \sin 2x$$

$$\begin{aligned}
 Q &= \int \frac{f_1 x}{\omega} dx \\
 &= \int \frac{\cos 2x + \tan 2x}{2} dx \\
 &= \frac{1}{2} \int \cos 2x + \frac{\sin 2x}{\cos 2x} dx \\
 &= \frac{1}{2} \int \sin 2x dx = \frac{1}{2} \left[ -\frac{\cos 2x}{2} \right]
 \end{aligned}$$

$$Q = - \cos 2x$$

$$PI = Pf_1 + Qf_2$$

$$\begin{aligned}
 &= \left[ \log |\sec 2x + \tan 2x| + \sin 2x \right] \cos 2x \\
 &\quad - \cos 2x \sin 2x
 \end{aligned}$$

$$PI = - \log |\sec 2x + \tan 2x| \cos 2x$$

$$\therefore y = CF + PI = C_1 \cos 2x + C_2 \sin 2x - \log |\sec 2x + \tan 2x| \cos 2x$$

