

Cauchy's Residue theorem:

[If $f(z)$ is analytic at all points inside and on a simple closed curve c except at a finite number of points $z_1, z_2, z_3, \dots, z_n$ inside c , then]


$$\oint_C f(z) dz = 2\pi i [\text{Sum of residues of } f(z) \text{ at } z_1, z_2, \dots, z_n]$$

Proof:

{ Given that $f(z)$ is not analytic

only at z_1, z_2, \dots, z_n .

Draw the non-intersecting small circles C_1, C_2, \dots, C_n with centres at z_1, z_2, \dots, z_n & radii r_1, r_2, \dots, r_n lying wholly inside c .



Then $f(z)$ is analytic in the region.

between c and c_1, c_2, \dots, c_n .

$$\int_C f(z) dz = \int_{c_1} f(z) dz + \int_{c_2} f(z) dz + \dots + \int_{c_n} f(z) dz \rightarrow ①$$

Now z_1, z_2, \dots, z_n are the singular points of $f(z)$.

$\therefore \{\text{Res } f(z)\}_{z=z_i}$ = the coef of $\frac{1}{z-z_i}$ in the Laurent's series of $f(z)$ about

$z=z_i$ (by defn of residues)

$$= b_1 = \frac{1}{2\pi i} \int_{c_i} \frac{f(z)}{(z-z_i)^{1-1}} dz$$

$$[\text{since } b_n = \frac{1}{2\pi i} \int_{c_i} \frac{f(z)}{(z-z_i)^{1-n}} dz]$$

$$= \frac{1}{2\pi i} \int_{c_i} \frac{f(z)}{z-z_i} dz$$

Since $\int_C f(z) dz$ is no bias
odd, obtain b_1 and b_n to odd part of $f(z)$.

$$\Rightarrow \int_C f(z) dz = 2\pi i \{\text{Res } f(z)\}_{z=z_i} \rightarrow ②$$

From ① & ②,

$$\begin{aligned} \int_C f(z) dz &= 2\pi i \{\text{Res } f(z)\}_{z=z_1} + 2\pi i \{\text{Res } f(z)\}_{z=z_2} \\ &\quad + \dots + 2\pi i \{\text{Res } f(z)\}_{z=z_n} \\ &= 2\pi i \left[\{\text{Res } f(z)\}_{z=z_1} + \{\text{Res } f(z)\}_{z=z_2} \right. \\ &\quad \left. + \dots + \{\text{Res } f(z)\}_{z=z_n} \right] \end{aligned}$$

$= 2\pi i \{ \text{Sum of residues of } f(z) \text{ at } z = z_1, z_2, \dots, z_n \}$

$$= 2\pi i \sum R_i$$

Q Evaluate $\int_C \frac{e^z}{(z+2)(z+1)^2} dz$ where C is the circle $|z| = 3$.

$$|z| = 3.$$

Soln:

$$\text{Let } f(z) = \frac{e^z}{(z+2)(z+1)^2}$$

The poles of $f(z)$ are given by,

$$(z+2)(z+1)^2 = 0 \Rightarrow z = -1, -2$$

$\Rightarrow z = -1$ is a pole of order 2.

$\Rightarrow z = -2$ is a pole of order 1.

$$\text{Given } |z| = 3$$

$z = -1 \Rightarrow |z| = 1 < 3$, lies inside C

$z = -2 \Rightarrow |z| = 2 > 3$, lies outside C

$$\{ \text{Res } f(z) \}_{z=-2} = \lim_{z \rightarrow -2} \frac{(z+2)}{(z+1)^2} \cdot \frac{e^z}{(z+2)(z+1)^2}$$

$$\frac{e^{-2}}{(-2+1)^2} = \frac{e^{-2}}{1} = e^{-2}$$

$$\{ \text{Res } f(z) \}_{z=-1} = \lim_{z \rightarrow -1} \frac{1}{2!} \frac{d^2}{dz^2} \left[\frac{e^z}{(z+1)^2 (z+2)} \right]$$

$$= \lim_{z \rightarrow -1} \frac{d}{dz} \frac{e^z}{(z+2)} = \frac{e^{-1}}{(-1+2)^2}$$

$$= \lim_{z \rightarrow -1} \frac{-e^{-1}}{(z+2)^2} = \frac{-e^{-1}}{(-1+2)^2} = -e^{-2}$$

∴ By Cauchy's residue theorem,

$\int_C f(z) dz = 2\pi i$ [Sum of the residues of $f(z)$ at the poles which lie inside C]

$$\int_C \frac{e^z dz}{(z+2)(z+1)^2} = 2\pi i \frac{(e^2 - e^2)}{(1+\infty)(2+\infty)} = 0.$$

(2) Evaluate $\int_C \frac{4-3z}{z^2(z-1)(z-2)} dz$ where C is the circle $|z| = 3/2$.

Soln:

$$\text{Let } f(z) = \frac{4-3z}{z(z-1)(z-2)}$$

The poles of $f(z)$ are,

$$z(z-1)(z-2) = 0$$

$z=0, z=1, z=2$ are poles of order 1.

$$z=0 \Rightarrow |z| = 0 < 3/2 \text{ lies inside } C$$

$$z=1 \Rightarrow |z| = 1 < 3/2 \text{ lies inside } C$$

$$z=2 \Rightarrow |z| = 2 > 3/2 \text{ lies outside } C$$

$$\{\text{Res } f(z)\}_{z=0} = \lim_{z \rightarrow 0} (z-0) \frac{4-3z}{z(z-1)(z-2)} = \frac{4}{(-1)(-2)} = 2$$

$$\{\text{Res } f(z)\}_{z=1} = \lim_{z \rightarrow 1} (z-1) \frac{4-3z}{z(z-1)(z-2)} = \frac{1}{(2)(-1)} = -1$$

$$\frac{1}{-1} = -1$$

$\text{Res}_{z=2} f(z) \neq 0$ (lies outside c)

By Cauchy's residue theorem,

$\int_C f(z) dz = 2\pi i$ [Sum of residues of $f(z)$ at the poles which lie inside C]

$$\int_{C(0, R)} f(z) dz = 2\pi i (2 - 1) \text{ (one pole at } z = 1 \text{ and none at } z = 2)$$

$$\text{and } \lim_{R \rightarrow \infty} \int_{C(0, R)} f(z) dz = 2\pi i \text{ (one pole at } z = 1 \text{ and none at } z = 2)$$