INVERSE LAPLACE TRANSFORM

Defn: If the Laplace Transform of $f(t)$ is $F(s)$ (ie) $L[f(t)] = F(s)$ Then $f(t)$ is called an Inverse Laplace Transform of $F(s)$ and is written as $f(t) = L^{-1}[F(s)]$ where L^{-1} is called the inverse Laplace transformation operator.

Table of ILT:

$L[f(t)] = F(s)$	$L^{-1}[F(s)] = f(t)$
1) $L[1] = \frac{1}{s}$	$\Rightarrow L^{-1}\left[\frac{1}{s}\right] = 1$
2) $L[t] = \frac{1}{s^2}$	$\Rightarrow L^{-1}\left[\frac{1}{s^2}\right] = t$
3) $L[t^n] = \frac{n!}{s^{n+1}}$	$\Rightarrow L^{-1}\left[\frac{n!}{s^{n+1}}\right] = t^n$
4) $L[e^{at}] = \frac{1}{s-a}$	$\Rightarrow L^{-1}\left[\frac{1}{s-a}\right] = e^{at}$
5) $L[e^{-at}] = \frac{1}{s+a}$	$\Rightarrow L^{-1}\left[\frac{1}{s+a}\right] = e^{-at}$
6) $L[\sin at] = \frac{a}{s^2+a^2}$	$\Rightarrow L^{-1}\left[\frac{a}{s^2+a^2}\right] = \sin at$
7) $L\left[\frac{\sin at}{a}\right] = \frac{1}{s^2+a^2}$	$\Rightarrow L^{-1}\left[\frac{1}{s^2+a^2}\right] = \frac{\sin at}{a}$

8) $L[\cos at] = \frac{s}{s^2+a^2} \Rightarrow L^{-1}\left[\frac{s}{s^2+a^2}\right] = \cos at$

9) $L[\sin hat] = \frac{a}{s^2-a^2} \Rightarrow L^{-1}\left[\frac{a}{s^2-a^2}\right] = \sin hat$

10) $L[\cos hat] = \frac{s}{s^2-a^2} \Rightarrow L^{-1}\left[\frac{s}{s^2-a^2}\right] = \cos hat$

11) $L[\delta(t)] = 1 \Rightarrow L^{-1}[1] = \delta(t)$